

Behavioral Counseling After Screening for Alcohol Misuse in Primary Care: A Systematic Review and Meta-analysis for the U.S. Preventive Services Task Force

Daniel E. Jonas, MD, MPH; James C. Garbutt, MD; Halle R. Amick, MSPH; Janice M. Brown, PhD; Kimberly A. Brownley, PhD; Carol L. Council, MSPH; Anthony J. Viera, MD, MPH; Tania M. Wilkins, MS; Cody J. Schwartz, MPH; Emily M. Richmond, MPH; John Yeatts, MPH; Tammeka Swinson Evans, MOP; Sally D. Wood, BA; and Russell P. Harris, MD, MPH

Background: Alcohol misuse, which includes the full spectrum from risky drinking to alcohol dependence, is a leading cause of preventable death in the United States.

Purpose: To evaluate the benefits and harms of behavioral counseling interventions for adolescents and adults who misuse alcohol.

Data Sources: MEDLINE, EMBASE, the Cochrane Library, CINAHL, PsycINFO, International Pharmaceutical Abstracts, and reference lists of published literature (January 1985 through January 2012, limited to English-language articles).

Study Selection: Controlled trials at least 6 months' duration that enrolled persons with alcohol misuse identified by screening in primary care settings and evaluated behavioral counseling interventions.

Data Extraction: One reviewer extracted data and a second checked accuracy. Two independent reviewers assigned quality ratings and graded the strength of the evidence.

Data Synthesis: The 23 included trials generally excluded persons with alcohol dependence. The best evidence was for brief (10- to 15-minute) multicontact interventions. Among adults receiving behavioral interventions, consumption decreased by 3.6 drinks per week from baseline (weighted mean difference, 3.6 drinks/wk [95% CI, 2.4 to 4.8 drinks/wk]; 10 trials; 4332 participants), 12%

fewer adults reported heavy drinking episodes (risk difference, 0.12 [CI, 0.07 to 0.16]; 7 trials; 2737 participants), and 11% more adults reported drinking less than the recommended limits (risk difference, 0.11 [CI, 0.08 to 0.13]; 9 trials; 5973 participants) over 12 months compared with control participants (moderate strength of evidence). Evidence was insufficient to draw conclusions about accidents, injuries, or alcohol-related liver problems. Trials enrolling young adults or college students showed reduced consumption and fewer heavy drinking episodes (moderate strength of evidence). Little or no evidence of harms was found.

Limitations: Results may be biased to the null because the behavior of control participants could have been affected by alcohol misuse assessments. In addition, evidence is probably inapplicable to persons with alcohol dependence and selective reporting may have occurred.

Conclusion: Behavioral counseling interventions improve behavioral outcomes for adults with risky drinking.

Primary Funding Source: Agency for Healthcare Research and Quality.

Ann Intern Med. 2012;157:645-654.

www.annals.org

For author affiliations, see end of text.

This article was published at www.annals.org on 25 September 2012.

Alcohol misuse, which includes the full spectrum from risky or hazardous drinking to alcohol dependence (1–3), is associated with numerous health and social problems and more than 85 000 deaths per year in the United States (4, 5). Alcohol misuse is the third leading cause of preventable death in the United States, after tobacco use and being overweight (6). It contributes to hypertension, cirrhosis, gastritis, gastric ulcers, pancreatitis, breast cancer, neuropathy, cardiomyopathy, anemia, osteoporosis, cognitive impairment, depression, insomnia, anxiety, suicide, injury, and violence (7–9). The definitions of the spectrum of alcohol misuse (that is, unhealthy alcohol use [1]) continue to evolve. For this review, we use the definitions in Table 1 (5, 10–12).

About 30% of the U.S. population misuse alcohol, with most engaging in what is considered risky drinking (1). Recent U.S.-based data (13) revealed that 21.3% of primary care patients reported risky drinking.

Cross-sectional and cohort studies have consistently related high average alcohol consumption and heavy per-occasion use to short- or long-term health consequences (14, 15). A meta-analysis examining the association be-

tween all-cause mortality and average alcohol consumption (16) found that men who drank an average of at least 4 drinks per day and women who drank an average of at least 2 drinks per day had increased mortality relative to nondrinkers. The National Institute on Alcohol Abuse and Alcoholism has proposed guidelines (17) to limit the risks for drinking-related consequences. The maximum recommended consumption is 3 or fewer standard drinks per day (≤ 7 drinks/wk) for adult women and anyone older than 65 years, and 4 or fewer standard drinks per day (≤ 14 drinks/week) for men (15, 17, 18). These guidelines do not apply to persons for whom alcohol intake is contraindicated, such as pregnant women, persons with alcohol dependence or medical conditions that can be worsened by drinking, or those receiving medications that interact with alcohol.

Behavioral counseling interventions include the range of personal counseling and related behavior-change interventions that are used to help patients change health-related behaviors (19). “Counseling” here denotes a cooperative method of work that demands active participation from both patient and clinician and aims to facilitate the

Table 1. Definitions of the Spectrum of Alcohol Misuse

| Term (Reference) | Definition |
|---|--|
| Risky or hazardous use (5) | Consumption of alcohol above recommended daily, weekly, or per-occasion amounts Consumption levels that increase the risk for health consequences |
| Harmful use (10, 11) | A pattern of drinking that is already causing damage to health; the damage may be either physical (e.g., liver damage from chronic drinking) or mental (e.g., depressive episodes secondary to drinking) |
| Alcohol abuse (12) | A maladaptive pattern of alcohol use leading to clinically significant impairment or distress, as manifested by ≥ 1 of the following within a 12-mo period: Recurrent alcohol use resulting in a failure to fulfill major obligations at work, school, or home (e.g., repeated absences or poor work performance related to alcohol use; alcohol-related absences, suspensions, or expulsions from school; or neglect of children or household) Recurrent alcohol use in situations in which it is physically hazardous (e.g., driving an automobile or operating a machine) Recurrent alcohol-related legal problems (e.g., arrests for alcohol-related disorderly conduct) Continued use despite persistent or recurrent social or interpersonal problems caused or exacerbated by the effects of alcohol (e.g., arguments with spouse about consequences of intoxication or physical fights) The symptoms have never met the criteria for alcohol dependence |
| Alcohol dependence (alcoholism, alcohol addiction) (12) | A maladaptive pattern of alcohol use leading to clinically significant impairment or distress, as manifested by ≥ 3 of the following at any time in the same 12-mo period: Tolerance, as defined by either of the following: A need for markedly increased amounts of alcohol to achieve intoxication or desired effect Markedly diminished effect with continued use of the same amount of alcohol Withdrawal, as manifested by either of the following: The characteristic withdrawal syndrome for alcohol Alcohol (or a closely related drug) is taken to relieve or avoid withdrawal symptoms Alcohol is often taken in larger amounts or over a longer period than was intended A persistent desire or unsuccessful efforts to cut down or control alcohol use A great deal of time is spent in activities necessary to obtain alcohol, use alcohol, or recover from its effects Important social, occupational, or recreational activities are given up or reduced because of alcohol use Use continues despite knowledge of a persistent or recurrent physical or psychological problem that is likely to have been caused or exacerbated by alcohol (e.g., continued drinking despite recognition that an ulcer was made worse by alcohol consumption) |

patient’s independent initiative (19). The goal of behavioral interventions for alcohol misuse is to eliminate risky drinking practices (for example, by encouraging fewer drinks per occasion or not drinking before driving) rather than to achieve abstinence.

For the Effective Health Care Program of the Agency for Healthcare Research and Quality (AHRQ) and to assist the U.S. Preventive Services Task Force (USPSTF) in updating its 2004 recommendation statement (20), we conducted a systematic review and meta-analysis of the effectiveness of screening followed by behavioral counseling, with or without referral, for alcohol misuse in primary care settings (21). The full report (21) addressed 7 questions (Appendix Table 1, available at www.annals.org).

METHODS

We developed and followed a standard protocol. A technical report that details methods and includes search strategies and additional evidence tables is available at www.effectivehealthcare.ahrq.gov/reports/final.cfm.

Key Questions and Analytic Framework

The USPSTF and the AHRQ determined the focus of this review. Investigators developed key questions and created an analytic framework that incorporated the key questions and outlined patient populations, interventions, comparators, outcomes (including adverse effects), and settings (Appendix Figure 1, available at www.annals.org). This

report focuses on the key questions related to benefits and harms of behavioral interventions.

Data Sources and Searches

We searched MEDLINE, EMBASE, the Cochrane Library, CINAHL, PsycINFO, and the International Pharmaceutical Abstracts from 1 January 1985 to 31 January 2012, limited to English-language articles. The start date was selected on the basis of the earliest publication date found in previous reviews and expert opinion. We used Medical Subject Headings as search terms when available and keywords when appropriate, focusing on terms to describe relevant populations, screening, and behavioral interventions.

Study Selection

We developed inclusion and exclusion criteria with respect to populations, interventions, comparators, outcomes, timing, settings, and study designs (22, 23). For the question related to behavioral interventions, we included randomized, controlled trials of at least 6 months’ duration that enrolled adults or adolescents with alcohol misuse identified by screening in primary care settings and that evaluated whether a counseling intervention improved behavioral or health outcomes.

Two investigators independently reviewed titles and abstracts, and then another 2 investigators independently reviewed the full text of all articles marked for possible inclusion during the initial review to determine final inclu-

sion or exclusion. Disagreements were resolved with an experienced team member.

Data Extraction and Quality Assessment

We designed and used structured forms to extract pertinent information from each article, including information about the methods and populations, interventions, comparators, outcomes, timing, settings, and study designs. All data extractions were reviewed for completeness and accuracy by a second team member.

We assessed the quality (internal validity) of studies using predefined criteria based on those developed by the USPSTF (ratings of good, fair, or poor) (24) and the University of York Centre for Reviews and Dissemination (25). These included assessment of the adequacy of randomization, allocation concealment, similarity of groups at baseline, masking, attrition, and whether intention-to-treat analysis was used. Two independent reviewers assigned quality ratings for each study. Disagreements were resolved by an experienced member of the team.

Data Synthesis and Analysis

We stratified evidence by population (adults, older adults, young adults or college students, and pregnant women). Quantitative analyses were conducted of outcomes reported by a sufficient number of studies that were homogeneous enough to justify combining their results. We used random-effects models. For the outcome of alcohol consumption, the effect measure was mean difference between the intervention and control groups for change from baseline in drinks per week. The percentages of patients who had episodes of heavy drinking and those who achieved recommended drinking limits were compared (between intervention and control groups) with a risk difference. Because follow-up periods varied, the analysis for all-cause mortality was based on deaths per person-year and the comparison between intervention and control groups was calculated as a risk ratio. Analyses were conducted by using Comprehensive Meta Analysis, version 2.2.055 (Bio-Stat, Englewood, New Jersey).

We used subgroup analyses to explore whether results differed by intensity, sex, country, deliverer of the intervention, or setting. The chi-square and I^2 statistics were calculated to assess heterogeneity in effects between studies (26, 27). When quantitative analyses were not appropriate (for example, because of heterogeneity, insufficient number of similar studies, or insufficient or varied outcome reporting), we synthesized the data qualitatively.

To assess the differential effects of using more or less time and single or multiple contacts, we grouped interventions by intensity of counseling, as measured by the duration and number of contacts: very brief (≤ 5 minutes, single-contact), brief (6 to 15 minutes, single-contact), extended (> 15 minutes, single-contact), brief multicontact (each contact ≤ 15 minutes), or extended multicontact (some contacts > 15 minutes).

We then graded the strength of evidence (SOE) as high, moderate, low, or insufficient on the basis of the guidance established for the Evidence-based Practice Center Program (Appendix Table 2, available at www.annals.org) (28). Two reviewers assessed each domain for each key outcome, and differences were resolved by consensus.

Role of the Funding Source

This review was funded by AHRQ. Staff of AHRQ and members of the USPSTF participated in developing the scope of the work and reviewed draft manuscripts. Approval from AHRQ for copyright assertion was required before the manuscript could be submitted for publication, but the authors are solely responsible for the content and the decision to submit it for publication.

RESULTS

We included 38 articles reporting on 23 randomized, controlled trials (Appendix Figure 2, available at www.annals.org). Sample sizes ranged from 72 to 1559, and study durations ranged from 6 to 48 months (Appendix Table 3, available at www.annals.org). Eleven studies were done solely in the United States, 2 focused on older adults, 5 focused on young adults or college students, and 1 enrolled pregnant women. We identified no studies of adolescents.

Fourteen of the interventions (29–50) were delivered by a primary care physician alone or with a health educator or nurse. Three (51–54) were delivered by a nurse or physician assistant, 1 by a psychologist (55–57), 2 by a researcher (58–62), and 1 by unspecified interventionists (63). Two interventions in college students (64–66) were conducted via a computer. Most trials tested brief multi-contact interventions (31–34, 42, 46, 50, 51, 53, 64, 65) or brief interventions (29, 49, 52, 58, 60, 62–66); fewer tested very brief (45, 63), extended (30), or extended multi-contact interventions (38, 45, 48, 55, 60). Interventions were heterogeneous and included various counseling approaches, such as brief advice, feedback, or motivational interviews, and cognitive behavioral strategies, such as self-completed action plans, written health education or self-help materials, drinking diaries, or problem-solving exercises to complete at home (Appendix Table 4, available at www.annals.org). Most comparator groups received screening or assessment followed by usual care or by provision of a general health pamphlet. A few studies included additional components in comparator groups that could have biased results toward the null, such as recording screening or assessment results on the chart (45) or forwarding them to physicians (60), advice from nurses on reducing drinking and a leaflet with benchmark alcohol guides (52), a pamphlet on the health effects of alcohol consumption (64–66), or a booklet about preventing alcohol problems (48). We summarize the main findings by population and outcome and report the SOE for each.

Table 2. Effectiveness and Strength of Evidence of Behavioral Interventions Compared With Controls for Improving Intermediate Outcomes, by Population*

| Population | Mean Consumption† | | Heavy Drinking Episodes‡ | | Recommended Drinking Limits | |
|-------------------------------|---|----------------------|---|----------------------|---|----------------------|
| | Results | Strength of Evidence | Results | Strength of Evidence | Results | Strength of Evidence |
| Adults | 3.6 fewer drinks/wk (95% CI, 2.4–4.8 drinks/wk); 10 trials; 4332 participants | Moderate | 12% fewer participants reported heavy drinking episodes (95% CI, 7%–16%); 7 trials; 2737 participants | Moderate | Achieved by 11% more participants (95% CI, 8%–13%); 9 trials; 5973 participants | Moderate |
| Older adults | 1.7 fewer drinks/wk (95% CI, 0.6–2.8 drinks/wk); 2 trials; 776 participants | Moderate | – | Insufficient | Achieved by 9% more participants (95% CI, 2%–16%); 2 trials; 776 participants | Low |
| Young adults/college students | 1.7 fewer drinks/wk (95% CI, 0.7–2.6 drinks/wk); 3 trials; 1421 participants | Moderate§ | 0.9 fewer heavy drinking day per month (95% CI, 0.3–1.5 fewer); 3 trials; 1448 participants | Moderate§ | – | Insufficient |
| Pregnant women | Data from 1 study found no difference | Low | – | Insufficient | – | Insufficient |
| Adolescents | – | Insufficient | – | Insufficient | – | Insufficient |

* All outcomes are at 12 mo unless indicated. All percentages are absolute risk differences from our meta-analyses.
 † Baseline consumption, adults: mean, about 23 drinks/wk; median, about 19 drinks/wk; range, 8–62 drinks/wk (data from 16 trials). Baseline consumption, older adults: 15.2–16.6 drinks/wk (data from 2 trials). Baseline consumption, young adults/college students: mean, about 15 drinks/wk; median, about 17 drinks/wk; range, 8–18 drinks/wk (2 of the 5 trials did not report).
 ‡ Generally defined as consumption of ≥5 standard drinks for men and ≥4 for women.
 § These data are 6-mo outcomes. For consumption for young adults, we could not calculate pooled point estimates for 12-mo data, but the range of reduction was 1.2–4.1 drinks/wk at 12 mo (moderate strength); for heavy drinking days for young adults, differences were not statistically significant at 12 mo (low strength).
 || Baseline heavy drinking days for young adults/college students, about 6–7 d over the past month.

Screening

We found no studies meeting inclusion criteria that randomly assigned participants, practices, or providers to screening and a comparator (no studies addressing questions 1 or 3) (Appendix Table 1). We found adequate evidence that several screening instruments can detect alcohol misuse in adults with acceptable sensitivity and specificity (21). The full technical report includes additional details about the accuracy of screening tests.

Effectiveness for Improving Intermediate Outcomes

Table 2 summarizes the results of meta-analyses for consumption, heavy drinking, and recommended drinking limits, by population. The Figure shows the forest plots for 12-month outcomes from our meta-analyses for adults. Overall, evidence supports the effectiveness of behavioral interventions for improving several intermediate outcomes for adults, older adults, and young adults or college students. For pregnant women, the included study (250 participants) (30) did not provide evidence of effectiveness for improving intermediate outcomes over 6 months or longer (low or insufficient SOE, depending on the outcome). Subgroup analyses identified no significant differences between men and women. Brief multicontact interventions had the best evidence of effectiveness across populations and outcomes and had follow-up data spanning several years. Meta-analyses of studies in adults found very brief and brief single-contact interventions to be ineffective for some outcomes and less effective than brief multicontact interventions for others.

Effectiveness for Reducing Morbidity, Reducing Mortality, or Changing Other Outcomes

Table 3 summarizes results, by population. Our meta-analyses found no statistically significant reduction in all-cause mortality for adults (rate ratio, 0.64 [95% CI, 0.24 to 1.7]; 4 trials) or for all age groups combined (rate ratio, 0.52 [CI, 0.22 to 1.2]; 6 trials). Point estimates trended toward favoring interventions, but few studies reported mortality and few long-term data were available. No studies that enrolled pregnant women and reported these outcomes were found (insufficient SOE).

Potential Adverse Effects

We found no evidence of direct harms, aside from opportunity costs associated with interventions, which ranged from 5 minutes to 2 hours dispersed over several in-person or telephone visits (moderate SOE). We searched for evidence of potential adverse effects, such as illegal substance use, increased smoking, anxiety, stigma, labeling, discrimination, or interference with the physician–patient relationship. We found no evidence for most of these potential harms and very limited evidence reporting no difference between groups for smoking rates and anxiety (low SOE). Other than the results for opportunity costs, our results are limited by the few trials that reported any information; 5 of 23 reported smoking (29, 33, 34, 39, 41, 49, 50), and 2 reported anxiety (29, 49).

Health Care System Influences


Where the study was conducted (United States vs. non–United States) had no impact on the effectiveness of

Figure. Forest plots for alcohol consumption, heavy drinking, and achieving recommended drinking limits for groups receiving behavioral counseling interventions compared with control groups.

Change in Alcohol Consumption From Baseline to 12 mo (drinks/wk)

| Study (Reference) | BCI Subgroup | Difference in Means (95% CI) | P Value |
|----------------------------------|------------------------|------------------------------|---------|
| Richmond et al (45) | Very brief | 2.700 (−5.212 to 10.612) | 0.50 |
| Subgroup total | | 2.700 (−5.212 to 10.612) | 0.50 |
| Anderson and Scott (29) | Brief | −4.740 (−9.544 to 0.064) | 0.053 |
| Scott and Anderson (49) | Brief | −1.600 (−8.227 to 5.027) | 0.64 |
| Lock et al (52) | Brief | −0.190 (−8.935 to 8.555) | 0.97 |
| ELM trial (59–61) | Brief | −4.430 (−8.545 to −0.315) | 0.035 |
| Subgroup total | | −3.660 (−6.349 to −0.970) | 0.008 |
| Project TrEAT (33, 35–37, 39) | Brief, multicontact | −4.180 (−5.887 to −2.473) | 0.000 |
| Project Health (42–44) | Brief, multicontact | −2.700 (−5.156 to −0.244) | 0.031 |
| Rubio et al (46) | Brief, multicontact | −3.560 (−4.898 to −2.222) | 0.000 |
| Wallace et al (50) (men) | Brief, multicontact | −10.100 (−14.400 to −5.800) | 0.000 |
| Wallace et al (50) (women) | Brief, multicontact | −5.200 (−10.252 to −0.148) | 0.044 |
| Subgroup total | | −4.407 (−6.084 to −2.730) | 0.000 |
| Richmond et al (45) | Extended, multicontact | −2.200 (−11.331 to 6.931) | 0.64 |
| ELM trial (59–61) | Extended, multicontact | −1.811 (−5.182 to 1.560) | 0.29 |
| SIP trial (55–57) (full care) | Extended, multicontact | −3.420 (−7.826 to 0.986) | 0.128 |
| SIP trial (55–57) (stepped care) | Extended, multicontact | −3.010 (−7.430 to 1.410) | 0.182 |
| Subgroup total | | −2.546 (−4.767 to −0.325) | 0.025 |
| Total | | −3.573 (−4.758 to −2.389) | 0.000 |


Heterogeneity statistics: $Q = 15.066$; $P = 0.303$; $I^2 = 13.714$



Achieved Recommended Drinking Limit at 12 mo

| Study (Reference) | BCI Subgroup | Risk Difference (95% CI) | P Value |
|---------------------------------------|---------------------|--------------------------|---------|
| WHO BISG (63) (men) | Very brief | 0.080 (0.012 to 0.148) | 0.021 |
| WHO BISG (63) (women) | Very brief | 0.080 (−0.058 to 0.218) | 0.26 |
| Subgroup total | | 0.080 (0.019 to 0.141) | 0.010 |
| WHO BISG (63) (men) | Brief | 0.080 (0.015 to 0.145) | 0.015 |
| WHO BISG (63) (women) | Brief | 0.040 (−0.099 to 0.179) | 0.57 |
| Anderson and Scott (29) | Brief | 0.130 (0.032 to 0.228) | 0.009 |
| Senft et al (62); Freeborn et al (58) | Brief | 0.070 (−0.003 to 0.143) | 0.060 |
| Scott and Anderson (49) | Brief | 0.010 (−0.195 to 0.215) | 0.92 |
| Subgroup total | | 0.079 (0.039 to 0.120) | 0.000 |
| Curry et al (31) | Brief, multicontact | 0.140 (0.029 to 0.251) | 0.013 |
| Project TrEAT (33, 35–37, 39) | Brief, multicontact | 0.134 (0.072 to 0.196) | 0.000 |
| Project Health (42–44) | Brief, multicontact | 0.050 (−0.043 to 0.143) | 0.29 |
| Rubio et al (46) | Brief, multicontact | 0.187 (0.117 to 0.256) | 0.000 |
| Wallace et al (50) (men) | Brief, multicontact | 0.182 (0.110 to 0.254) | 0.000 |
| Wallace et al (50) (women) | Brief, multicontact | 0.185 (0.070 to 0.300) | 0.002 |
| Subgroup total | | 0.149 (0.109 to 0.188) | 0.000 |
| Total | | 0.109 (0.083 to 0.134) | 0.000 |


Heterogeneity statistics: $Q = 17.366$; $P = 0.136$; $I^2 = 30.900$



No Heavy Drinking Episodes at 12 mo

| Study (Reference) | BCI Subgroup | Risk Difference (95% CI) | P Value |
|----------------------------------|------------------------|--------------------------|---------|
| Anderson and Scott (29) | Brief | 0.167 (0.023 to 0.311) | 0.023 |
| Scott and Anderson (49) | Brief | 0.030 (−0.128 to 0.188) | 0.71 |
| Subgroup total | | 0.102 (−0.032 to 0.236) | 0.134 |
| Curry et al (31) | Brief, multicontact | 0.050 (−0.033 to 0.133) | 0.24 |
| Project TrEAT (33, 35–37, 39) | Brief, multicontact | 0.141 (0.074 to 0.208) | 0.000 |
| Project Health (42–44) | Brief, multicontact | 0.060 (−0.033 to 0.153) | 0.20 |
| Rubio et al (46) | Brief, multicontact | 0.149 (0.080 to 0.218) | 0.000 |
| Subgroup total | | 0.106 (0.056 to 0.157) | 0.000 |
| SIP trial (55–57) (full care) | Extended, multicontact | 0.189 (0.020 to 0.358) | 0.029 |
| SIP trial (55–57) (stepped care) | Extended, multicontact | 0.193 (0.031 to 0.355) | 0.020 |
| Subgroup total | | 0.191 (0.074 to 0.308) | 0.001 |
| Total | | 0.118 (0.074 to 0.162) | 0.000 |

Heterogeneity statistics: $Q = 8.457$; $P = 0.294$; $I^2 = 17.232$



BCI = behavioral counseling intervention; ELM = Early Lifestyle Modification; SIP = Screening and Intervention in Primary Care; TrEAT = Trial for Early Alcohol Treatment; WHO BISG = World Health Organization Brief Intervention Study Group.

Table 3. Effectiveness and Strength of Evidence of Behavioral Interventions Compared With Controls for Improving Health, Utilization, and Other Outcomes, by Population*

| Outcomes | Adults | | Older Adults | Young Adults/College Students | |
|--------------------------------|---|----------------------|----------------------|--|----------------------|
| | Results | Strength of Evidence | Strength of Evidence | Results | Strength of Evidence |
| Health outcomes | | | | | |
| Mortality | Rate ratio, 0.64 (95% CI, 0.24 to 1.7); 4 trials; 2006 participants† | Low | Insufficient | 1 death reported in a control group | Insufficient |
| Alcohol-related accidents‡ | – | Insufficient | Insufficient | Fewer motor vehicle crashes with nonfatal injuries (9 vs. 20 crashes; $P < 0.05$) and fewer total motor vehicle events (114 vs. 149 events; $P < 0.05$) after 48 mo§ | Low |
| Alcohol-related liver problems | – | Insufficient | Insufficient | – | Insufficient |
| Utilization outcomes | | | | | |
| Hospitalization | Fewer hospital days in the past 6 mo at 6, 12, and 48 mo: 35 vs. 180 d, 91 vs. 146 d, and 420 vs. 664 d, respectively; all $P < 0.05$ § | Low | Insufficient | Fewer hospital days but no statistically significant difference (131 vs. 150 d; $P =$ not significant)§ | Low |
| Emergency visits | No statistically significant difference | Low | Insufficient | Fewer emergency department visits (103 vs. 177 visits; $P < 0.01$)§ | Low |
| Primary care visits | No significant difference (weighted mean difference, -0.14 visits [95% CI, -0.5 to 0.2 visits]; 4 trials; 946 participants) | Low | Insufficient | – | Insufficient |
| Other outcomes | | | | | |
| Academic problems | – | – | – | Fewer consequences related to academic role expectations: rate ratio between 0.70 and 0.80¶ | Moderate |
| Legal problems** | No significant difference over 48 mo for most legal problems but fewer controlled substance/liquor violations (2 vs. 11 violations; $P < 0.05$)§ | Low | Insufficient | No significant difference for most legal problems but fewer controlled substance/liquor violations (0 vs. 8 violations; $P < 0.01$)§ | Low |
| Quality of life | No difference found in 3 trials (353 participants) | Low | Insufficient | – | Insufficient |

* Data are reported for 12-mo outcomes unless otherwise noted.
 † A meta-analysis that combined all age groups (adults, older adults, and young adults/college students) also found no statistically significant reduction in mortality (rate ratio, 0.52 [95% CI, 0.22 to 1.2]; 6 trials; 2255 participants), although point estimates trended toward favoring behavioral interventions. Few trials reported mortality, additional studies would be needed to increase precision, and few long-term data are available.
 ‡ “Accidents” indicates motor vehicle events and injuries.
 § Data are from Project TrEAT (Trial for Early Alcohol Treatment) (33, 35, 36), the best available evidence. The data for young adults are from Project TrEAT subgroup analyses (226 participants).
 || Results trended in favor of the intervention group at 6, 12, and 48 mo: 47 vs. 70 visits ($P > 0.10$), 60 vs. 62 visits ($P > 0.10$), and 302 vs. 376 visits ($P < 0.10$), respectively (33, 35, 36).
 ¶ Evidence from 2 trials (576 and 104 participants) conducted in New Zealand.
 ** Includes assault, battery, or child abuse; resisting or obstructing an officer or disorderly conduct; criminal or property damage; theft or robbery; and other arrests.

interventions for consumption outcomes. Data showed a tendency toward greater reduction in consumption for interventions delivered in academic- or research-oriented settings than for those delivered in community-based settings (weighted mean difference, -5.0 drinks/wk [CI, -7.6 to -2.5 drinks/wk] vs. -3.2 drinks/wk [CI, -4.3 to -2.2 drinks/wk]; 3 vs. 7 trials). Interventions delivered mostly by primary care providers showed a tendency toward greater reduction in consumption than did those delivered primarily by research personnel (weighted mean difference, -4.0 drinks/wk [CI, -5.4 to -2.6 drinks/wk] vs. -3.0 drinks/wk [CI, -5.0 to -1.0 drinks/wk]; 7 vs. 2 trials). Our consumption meta-analysis included only 1 intervention delivered by a nurse (52), and the reduction was not statistically significant in that study (weighted mean difference, -0.2 drinks/wk [CI, -8.9 to 8.6 drinks/wk]). Two

other studies, each of which provided insufficient data for our consumption meta-analysis, reported benefits of interventions delivered primarily by nurses (51) or by nurses and physician assistants (53) for some consumption outcomes. In addition, 2 interventions conducted by computer reported some evidence of effectiveness for reduced consumption in college students (64–66).

DISCUSSION

We found no studies that directly addressed our overarching question (key question 1)—no studies randomly assigned patients, practices, or providers to screening and comparator groups and subsequently provided interventions for those with positive screening results. All of the

included studies randomly assigned patients after they had received positive screening results.

We found that behavioral counseling interventions improved drinking behavior outcomes (moderate SOE) and reduced hospital days (low SOE) for adults with risky drinking. For most health outcomes, available evidence either found no difference between intervention and control groups, such as for mortality (low SOE), or was insufficient to draw conclusions, such as for alcohol-related liver problems (insufficient SOE). Long-term outcomes from 2 studies (33, 35–37, 39, 42, 43) revealed that participants in the intervention groups maintained reductions in consumption or continued to reduce consumption, but differences between intervention and control groups were no longer statistically significant by 48 months. Studies identified delayed reduction in consumption in control groups that could reflect the natural history of alcohol consumption, the cumulative effect of follow-up with the health care system, differential attrition (if more participants lost to follow-up in the control group were risky drinkers), or (late) regression to the mean.

The evidence for effectiveness in adults is strongest for brief multicontact interventions. The effect sizes for these interventions were greater than those for other intensities (although CIs often overlapped). In addition, the best studies show that the effect of brief multicontact interventions remains for several years (35, 36, 43) and also show improvement for some utilization outcomes, such as fewer hospital days (35, 36) and costs (benefit–cost ratio of 39:1 over 48 months [CI, 5.4 to 72.5]) (36).

The brief multicontact interventions generally lasted 10 to 15 minutes per contact. All of the brief multicontact interventions in our meta-analyses of behavioral outcomes at 12 months were delivered by primary care providers, sometimes with additional intervention from a nurse or health educator. For example, the intervention in Project TrEAT (Trial for Early Alcohol Treatment) (33) included two 15-minute visits with a primary care provider 1 month apart and two 5-minute follow-up phone calls from a nurse 2 weeks after each visit. The intervention also included feedback about health behaviors, a review of problem drinking prevalence, a list of the adverse effects of alcohol, a worksheet on drinking cues, a drinking agreement or prescription, and drinking diary cards. Of note, 2 studies of brief multicontact interventions in adults, both of which provided insufficient data for our meta-analyses, reported benefits of interventions delivered primarily by nurses (51) or by nurses and physician assistants (53) for some consumption outcomes.

Evidence suggests that very brief interventions (up to 5 minutes, single-contact) and brief interventions (up to 15 minutes, single-contact) are less effective or ineffective, depending on the outcome. Although extended multicontact interventions seem to be effective for improving intermediate outcomes, we found no evidence that they are more effective than brief multicontact interventions.

The only included study that enrolled pregnant women (250 participants) (30) found no difference in reduced consumption between groups but did find higher rates of continued abstinence among women who were abstinent before the assessment in the intervention group than among those in the control group. Our searches identified other studies focusing on pregnant women that did not meet our inclusion criteria (67–84). Several took place in such settings as jails or specialized drug and alcohol treatment centers (75), and others lacked a control group or followed participants for fewer than 6 months (73, 84). Several of these studies reported benefits of interventions, including reduced consumption (73, 84), reduced risk for an alcohol-exposed pregnancy (75), higher rates of abstinence (79), and better fetal and newborn outcomes (higher birth weights and lengths and reduced fetal mortality rates [79]).

We have described several categories of alcohol misuse (such as risky or hazardous use and alcohol dependence). These categories are not all discrete (an individual may meet the definition for more than one). Included trials generally enrolled participants with risky or hazardous drinking, but the trials used varying terminology to describe the populations and often enrolled heterogeneous samples. Nevertheless, most investigators excluded participants with alcohol dependence or constructed their inclusion and exclusion criteria to substantially limit the number of such participants. Our best assessment is that our overall findings apply to risky or hazardous drinkers but not to persons with alcohol dependence. It is uncertain whether our findings apply to harmful drinkers or persons with alcohol abuse.

All interventions required support systems to provide screening; screening-related assessment; and in some cases, provider prompting. Screening assessments were often multistep processes that included interviews with research personnel that lasted up to 30 minutes. Less time would be required for screening and screening-related assessments in primary care practice. We estimate that 5 to 10 minutes would be required for persons who had positive screening results, with most of the time used to assess whether such persons have alcohol abuse or dependence (and should probably be referred for specialized treatment) as opposed to risky or hazardous drinking (for which behavioral counseling interventions in primary care may be effective). Nevertheless, support systems are probably required for effective screening and intervention. In addition, most interventions required training providers or staff.

It is unclear whether our findings apply to persons with certain comorbid conditions, and some researchers have suggested that brief behavioral interventions may be ineffective or less effective in people with comorbid psychiatric conditions. A subgroup analysis from a German study (56) found that brief interventions did not reduce drinking among 88 participants with comorbid anxiety or depression. Although most trials in our review did not exclude

persons with depression, anxiety, or chronic pain, it is unclear how many participants with these conditions were included in most trials.

A previous systematic review (85) found no evidence of efficacy for brief behavioral interventions in patients with alcohol dependence in primary care settings. Our review also found no such evidence. Included studies that enrolled more than 10% of participants with alcohol dependence reported interventions to be ineffective or less effective than studies that did not enroll alcohol-dependent participants.

Screening for alcohol misuse will inevitably identify some alcohol-dependent individuals; thus, providers and those making recommendations need information about whether effective interventions are available for alcohol dependence. If complete abstinence is used as an outcome, 15% to 35% of patients have been reported to achieve 1 year of sobriety after such treatment approaches (86) as pharmacotherapy, motivational enhancement therapy, cognitive behavioral therapy, 12-step facilitation, and therapy at alcoholism-treatment centers. Similar sobriety outcomes at 3 to 5 years or longer have been reported (9).

Our review has limitations. First, the scope of our review was limited to primary care settings. Second, most evidence involved self-report of alcohol use. Investigators in some trials verified self-reported use with other persons (such as family members). Self-report of alcohol use has been found to be accurate if collected carefully (87, 88). Third, the assessments conducted in the included trials could have concealed benefits of interventions (and biased results toward the null) by causing behavior changes. Control participants generally reduced alcohol consumption. Possible explanations include increased awareness of drinking, discussions with their provider about drinking that were prompted by the screening questions, receipt of some minimal intervention (control groups in the included studies often received some printed educational materials), or regression to the mean. A recent systematic review (89) concluded that answering questions on drinking in brief intervention trials seems to alter subsequent self-reported behavior, potentially generating bias by exposing nonintervention control groups to an integral component of the intervention. Finally, publication bias and selective reporting may be present.

In conclusion, behavioral counseling interventions improve intermediate outcomes, such as alcohol consumption, heavy drinking episodes, and drinking above recommended amounts (moderate SOE) and may reduce hospital days (low SOE) for adults with risky or hazardous drinking. For most health outcomes, available evidence found no difference between intervention and control groups, such as for mortality (low SOE), or was insufficient to draw conclusions about the effectiveness of behavioral interventions, such as for alcohol-related accidents or quality of life (insufficient SOE). Brief multicontact interven-

tions (about 10 to 15 minutes per contact) have the best evidence of effectiveness for adults.

From University of North Carolina at Chapel Hill and Cecil G. Sheps Center for Health Services Research, Chapel Hill, and Research Triangle Institute International, Research Triangle Park, North Carolina.

Disclaimer: The views expressed in this article do not represent and should not be construed to represent a determination or policy of AHRQ or the U.S. Department of Health and Human Services.

Grant Support: By AHRQ, contract 290-2007-10056-I.

Potential Conflicts of Interest: Disclosures can be viewed at www.acponline.org/authors/icmje/ConflictOfInterestForms.do?msNum=M11-3047.

Requests for Single Reprints: Daniel E. Jonas, MD, MPH, University of North Carolina at Chapel Hill, Department of Medicine, 5034 Old Clinic Building, CB 7110, Chapel Hill, NC 27599; e-mail, daniel_jonas@med.unc.edu.

Current author addresses and author contributions are available at www.annals.org.

References

1. Saitz R. Clinical practice. Unhealthy alcohol use. *N Engl J Med*. 2005;352:596-607. [PMID: 15703424]
2. U.S. Department of Veterans Affairs. AUDIT-C Frequently Asked Questions. Washington, DC: U.S. Department of Veterans Affairs; 2010. Accessed at www.queri.research.va.gov/tools/alcohol-misuse/alcohol-faqs.cfm on 8 June 2012.
3. Whitlock EP, Green CA, Polen MR. Behavioral Counseling Interventions in Primary Care to Reduce Risky/Harmful Alcohol Use. Systematic Evidence Review no. 30. Rockville, MD: Agency for Healthcare Research and Quality; 2004.
4. Mokdad AH, Marks JS, Stroup DF, Gerberding JL. Actual causes of death in the United States, 2000. *JAMA*. 2004;291:1238-45. [PMID: 15010446]
5. Whitlock EP, Polen MR, Green CA, Orleans T, Klein J; U.S. Preventive Services Task Force. Behavioral counseling interventions in primary care to reduce risky/harmful alcohol use by adults: a summary of the evidence for the U.S. Preventive Services Task Force. *Ann Intern Med*. 2004;140:557-68. [PMID: 15068985]
6. Centers for Disease Control and Prevention. FastStats: Alcohol Use. Atlanta: Centers for Disease Control and Prevention; 2012. Accessed at www.cdc.gov/nchs/fastats/alcohol.htm on 8 June 2012.
7. Cherpitel CJ, Ye Y. Alcohol-attributable fraction for injury in the U.S. general population: data from the 2005 National Alcohol Survey. *J Stud Alcohol Drugs*. 2008;69:535-8. [PMID: 18612569]
8. Corrao G, Bagnardi V, Zambon A, La Vecchia C. A meta-analysis of alcohol consumption and the risk of 15 diseases. *Prev Med*. 2004;38:613-9. [PMID: 15066364]
9. Schuckit MA. Alcohol-use disorders. *Lancet*. 2009;373:492-501. [PMID: 19168210]
10. Isaac M, Janca A, Sartorius N. ICD-10 Symptom Glossary for Mental Disorders. Geneva: World Health Organization; 1994.
11. Janca A, Ustun TB, van Drimmelen J, Dittmann V, Isaac M. ICD-10 Symptom Checklist for Mental Disorders, version 1.1. Geneva: World Health Organization; 1994.
12. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition. Washington, DC: American Psychiatric Assoc; 1994.
13. Vinson DC, Manning BK, Galliher JM, Dickinson LM, Pace WD, Turner BJ. Alcohol and sleep problems in primary care patients: a report from the AAFP National Research Network. *Ann Fam Med*. 2010;8:484-92. [PMID: 21060117]

14. Bondy SJ, Rehm J, Ashley MJ, Walsh G, Single E, Room R. Low-risk drinking guidelines: the scientific evidence. *Can J Public Health*. 1999;90:264-70. [PMID: 10489725]
15. Shalala DE. 10th Special Report to the U.S. Congress on Alcohol and Health: Highlights From Current Research: From the Secretary of Health and Human Services. Washington, DC: U.S. Department of Health and Human Services; 2000. Accessed at <http://pubs.niaaa.nih.gov/publications/10report/intro.pdf> on 8 June 2012.
16. Holman CD, English DR, Milne E, Winter MG. Meta-analysis of alcohol and all-cause mortality: a validation of NHMRC recommendations. *Med J Aust*. 1996;164:141-5. [PMID: 8628131]
17. National Institute on Alcohol Abuse and Alcoholism. Helping Patients Who Drink Too Much: A Clinician's Guide. Updated 2005 ed. Washington, DC: U.S. Department of Health and Human Services; 2005. Accessed at <http://pubs.niaaa.nih.gov/publications/practitioner/cliniciansguide2005/guide.pdf> on 8 June 2012.
18. Dawson DA, Grant BF, Li TK. Quantifying the risks associated with exceeding recommended drinking limits. *Alcohol Clin Exp Res*. 2005;29:902-8. [PMID: 15897737]
19. Whitlock EP, Orleans CT, Pender N, Allan J. Evaluating primary care behavioral counseling interventions: an evidence-based approach. *Am J Prev Med*. 2002;22:267-84. [PMID: 11988383]
20. U.S. Preventive Services Task Force. Screening and behavioral counseling interventions in primary care to reduce alcohol misuse: recommendation statement. *Ann Intern Med*. 2004;140:554-6. [PMID: 15068984]
21. Jonas DE, Garbutt JC, Brown JM, Amick HR, Brownley KA, Council CL, et al. Screening, Behavioral Counseling, and Referral in Primary Care to Reduce Alcohol Misuse. Comparative Effectiveness Review no. 64. AHRQ Publication no. 12-EHC055-EF. Rockville, MD: Agency for Healthcare Research and Quality; 2012.
22. Counsell C. Formulating questions and locating primary studies for inclusion in systematic reviews. *Ann Intern Med*. 1997;127:380-7. [PMID: 9273830]
23. Agency for Healthcare Research and Quality. AHRQ EHCP Methods Guide for Comparative Effectiveness Reviews of Medical Interventions. Rockville, MD: Agency for Healthcare Research and Quality; 2010.
24. Harris RP, Helfand M, Woolf SH, Lohr KN, Mulrow CD, Teutsch SM, et al; Methods Work Group, Third U.S. Preventive Services Task Force. Current methods of the US Preventive Services Task Force: a review of the process. *Am J Prev Med*. 2001;20:21-35. [PMID: 11306229]
25. Centre for Reviews and Dissemination. Systematic Reviews: CRD's Guidance for Undertaking Reviews in Healthcare. York, UK: Univ York; 2009.
26. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med*. 2002;21:1539-58. [PMID: 12111919]
27. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ*. 2003;327:557-60. [PMID: 12958120]
28. Owens DK, Lohr KN, Atkins D, Treadwell JR, Reston JT, Bass EB, et al. AHRQ series paper 5: grading the strength of a body of evidence when comparing medical interventions—Agency for Healthcare Research and Quality and the Effective Health-Care Program. *J Clin Epidemiol*. 2010;63:513-23. [PMID: 19595577]
29. Anderson P, Scott E. The effect of general practitioners' advice to heavy drinking men. *Br J Addict*. 1992;87:891-900. [PMID: 1525531]
30. Chang G, Wilkins-Haug L, Berman S, Goetz MA. Brief intervention for alcohol use in pregnancy: a randomized trial. *Addiction*. 1999;94:1499-508. [PMID: 10790902]
31. Curry SJ, Ludman EJ, Grothaus LC, Donovan D, Kim E. A randomized trial of a brief primary-care-based intervention for reducing at-risk drinking practices. *Health Psychol*. 2003;22:156-65. [PMID: 12683736]
32. Fleming MF, Balousek SL, Grossberg PM, Mundt MP, Brown D, Wiegel JR, et al. Brief physician advice for heavy drinking college students: a randomized controlled trial in college health clinics. *J Stud Alcohol Drugs*. 2010;71:23-31. [PMID: 20105410]
33. Fleming MF, Barry KL, Manwell LB, Johnson K, London R. Brief physician advice for problem alcohol drinkers. A randomized controlled trial in community-based primary care practices. *JAMA*. 1997;277:1039-45. [PMID: 9091691]
34. Fleming MF, Manwell LB, Barry KL, Adams W, Stauffacher EA. Brief physician advice for alcohol problems in older adults: a randomized community-based trial. *J Fam Pract*. 1999;48:378-84. [PMID: 10334615]
35. Fleming MF, Mundt MP, French MT, Manwell LB, Stauffacher EA, Barry KL. Benefit-cost analysis of brief physician advice with problem drinkers in primary care settings. *Med Care*. 2000;38:7-18. [PMID: 10630716]
36. Fleming MF, Mundt MP, French MT, Manwell LB, Stauffacher EA, Barry KL. Brief physician advice for problem drinkers: long-term efficacy and benefit-cost analysis. *Alcohol Clin Exp Res*. 2002;26:36-43. [PMID: 11821652]
37. Grossberg PM, Brown DD, Fleming MF. Brief physician advice for high-risk drinking among young adults. *Ann Fam Med*. 2004;2:474-80. [PMID: 15506584]
38. Lin JC, Karno MP, Tang L, Barry KL, Blow FC, Davis JW, et al. Do health educator telephone calls reduce at-risk drinking among older adults in primary care? *J Gen Intern Med*. 2010;25:334-9. [PMID: 20101471]
39. Manwell LB, Fleming MF, Mundt MP, Stauffacher EA, Barry KL. Treatment of problem alcohol use in women of childbearing age: results of a brief intervention trial. *Alcohol Clin Exp Res*. 2000;24:1517-24. [PMID: 11045860]
40. Moore AA, Blow FC, Hoffing M, Welgreen S, Davis JW, Lin JC, et al. Primary care-based intervention to reduce at-risk drinking in older adults: a randomized controlled trial. *Addiction*. 2011;106:111-20. [PMID: 21143686]
41. Mundt MP, French MT, Roebuck MC, Manwell LB, Barry KL. Brief physician advice for problem drinking among older adults: an economic analysis of costs and benefits. *J Stud Alcohol*. 2005;66:389-94. [PMID: 16047528]
42. Ockene JK, Adams A, Hurley TG, Wheeler EV, Hebert JR. Brief physician- and nurse practitioner-delivered counseling for high-risk drinkers: does it work? *Arch Intern Med*. 1999;159:2198-205. [PMID: 10527297]
43. Ockene JK, Reed GW, Reiff-Hekking S. Brief patient-centered clinician-delivered counseling for high-risk drinking: 4-year results. *Ann Behav Med*. 2009;37:335-42. [PMID: 19707840]
44. Reiff-Hekking S, Ockene JK, Hurley TG, Reed GW. Brief physician and nurse practitioner-delivered counseling for high-risk drinking. Results at 12-month follow-up. *J Gen Intern Med*. 2005;20:7-13. [PMID: 15693921]
45. Richmond R, Heather N, Wodak A, Kehoe L, Webster I. Controlled evaluation of a general practice-based brief intervention for excessive drinking. *Addiction*. 1995;90:119-32. [PMID: 7888970]
46. Rubio G, Jiménez-Arriero MA, Martínez I, Ponce G, Palomo T. Efficacy of physician-delivered brief counseling intervention for binge drinkers. *Am J Med*. 2010;123:72-8. [PMID: 20102995]
47. Saitz R, Horton NJ, Sullivan LM, Moskowitz MA, Samet JH. Addressing alcohol problems in primary care: a cluster randomized, controlled trial of a systems intervention. The Screening and Intervention in Primary Care (SIP) study. *Ann Intern Med*. 2003;138:372-82. [PMID: 12614089]
48. Schaus JF, Sole ML, McCoy TP, Mullett N, O'Brien MC. Alcohol screening and brief intervention in a college student health center: a randomized controlled trial. *J Stud Alcohol Drugs Suppl*. 2009;131-41. [PMID: 19538921]
49. Scott E, Anderson P. Randomized controlled trial of general practitioner intervention in women with excessive alcohol consumption. *Drug Alcohol Rev*. 1991;10:313-21. [PMID: 16818295]
50. Wallace P, Cutler S, Haines A. Randomised controlled trial of general practitioner intervention in patients with excessive alcohol consumption. *BMJ*. 1988;297:663-8. [PMID: 3052668]
51. Fleming MF, Lund MR, Wilton G, Landry M, Scheets D. The Healthy Moms Study: the efficacy of brief alcohol intervention in postpartum women. *Alcohol Clin Exp Res*. 2008;32:1600-6. [PMID: 18627361]
52. Lock CA, Kaner E, Heather N, Doughty J, Crawshaw A, McNamee P, et al. Effectiveness of nurse-led brief alcohol intervention: a cluster randomized controlled trial. *J Adv Nurs*. 2006;54:426-39. [PMID: 16671972]
53. Noknoy S, Rangsin R, Saengcharnchai P, Tantibhaedhyangkul U, McCambridge J. RCT of effectiveness of motivational enhancement therapy delivered by nurses for hazardous drinkers in primary care units in Thailand. *Alcohol Alcohol*. 2010;45:263-70. [PMID: 20236990]
54. Wilton G, Moberg DP, Fleming MF. The effect of brief alcohol intervention on postpartum depression. *MCN Am J Matern Child Nurs*. 2009;34:297-302. [PMID: 19713798]
55. Bischof G, Grothues JM, Reinhardt S, Meyer C, John U, Rumpf HJ. Evaluation of a telephone-based stepped care intervention for alcohol-related disorders: a randomized controlled trial. *Drug Alcohol Depend*. 2008;93:244-51. [PMID: 18054443]
56. Grothues JM, Bischof G, Reinhardt S, Meyer C, John U, Rumpf HJ. Effectiveness of brief alcohol interventions for general practice patients with problematic drinking behavior and comorbid anxiety or depressive disorders. *Drug Alcohol Depend*. 2008;94:214-20. [PMID: 18207336]

57. Reinhardt S, Bischof G, Grothues J, John U, Meyer C, Rumpf HJ. Gender differences in the efficacy of brief interventions with a stepped care approach in general practice patients with alcohol-related disorders. *Alcohol Alcohol*. 2008;43:334-40. [PMID: 18263901]
58. Freeborn DK, Polen MR, Hollis JF, Senft RA. Screening and brief intervention for hazardous drinking in an HMO: effects on medical care utilization. *J Behav Health Serv Res*. 2000;27:446-53. [PMID: 11070638]
59. Gordon AJ, Conigliaro J, Maisto SA, McNeil M, Kraemer KL, Kelley ME. Comparison of consumption effects of brief interventions for hazardous drinking elderly. *Subst Use Misuse*. 2003;38:1017-35. [PMID: 12901447]
60. Maisto SA, Conigliaro J, McNeil M, Kraemer K, Conigliaro RL, Kelley ME. Effects of two types of brief intervention and readiness to change on alcohol use in hazardous drinkers. *J Stud Alcohol*. 2001;62:605-14. [PMID: 11702799]
61. Maisto SA, Conigliaro J, McNeil M, Kraemer K, Kelley ME. The relationship between eligibility criteria for participation in alcohol brief intervention trials and other alcohol and health-related variables. *Am J Addict*. 2001;10:218-31. [PMID: 11579620]
62. Senft RA, Polen MR, Freeborn DK, Hollis JF. Brief intervention in a primary care setting for hazardous drinkers. *Am J Prev Med*. 1997;13:464-70. [PMID: 9415794]
63. WHO Brief Intervention Study Group. A cross-national trial of brief interventions with heavy drinkers. *Am J Public Health*. 1996;86:948-55. [PMID: 8669518]
64. Kypri K, Langley JD, Saunders JB, Cashell-Smith ML. Assessment may conceal therapeutic benefit: findings from a randomized controlled trial for hazardous drinking. *Addiction*. 2007;102:62-70. [PMID: 17207124]
65. Kypri K, Langley JD, Saunders JB, Cashell-Smith ML, Herbison P. Randomized controlled trial of web-based alcohol screening and brief intervention in primary care. *Arch Intern Med*. 2008;168:530-6. [PMID: 18332300]
66. Kypri K, Saunders JB, Williams SM, McGee RO, Langley JD, Cashell-Smith ML, et al. Web-based screening and brief intervention for hazardous drinking: a double-blind randomized controlled trial. *Addiction*. 2004;99:1410-7. [PMID: 15500594]
67. Armstrong MA, Kaskutas LA, Witbrodt J, Taillac CJ, Hung YY, Osejo VM, et al. Using drink size to talk about drinking during pregnancy: a randomized clinical trial of Early Start Plus. *Soc Work Health Care*. 2009;48:90-103. [PMID: 19197768]
68. Budd KW, Ross-Alaolmolki K, Zeller RA. Two prenatal alcohol use screening instruments compared with a physiologic measure. *J Obstet Gynecol Neonatal Nurs*. 2000;29:129-36. [PMID: 10750678]
69. Bull LB, Kvigne VL, Leonardson GR, Lacina L, Welty TK. Validation of a self-administered questionnaire to screen for prenatal alcohol use in Northern Plains Indian women. *Am J Prev Med*. 1999;16:240-3. [PMID: 10198664]
70. Chang G, Goetz MA, Wilkins-Haug L, Berman S. A brief intervention for prenatal alcohol use: an in-depth look. *J Subst Abuse Treat*. 2000;18:365-9. [PMID: 10812310]
71. Chang G, Goetz MA, Wilkins-Haug L, Berman S. Identifying prenatal alcohol use: screening instruments versus clinical predictors. *Am J Addict*. 1999;8:87-93. [PMID: 10365188]
72. Chang G, Wilkins-Haug L, Berman S, Goetz MA, Behr H, Hiley A. Alcohol use and pregnancy: improving identification. *Obstet Gynecol*. 1998;91:892-8. [PMID: 9610992]
73. Chang G, McNamara TK, Orav EJ, Koby D, Lavigne A, Ludman B, et al. Brief intervention for prenatal alcohol use: a randomized trial. *Obstet Gynecol*. 2005;105:991-8. [PMID: 15863535]
74. Chang G, McNamara TK, Orav EJ, Wilkins-Haug L. Brief intervention for prenatal alcohol use: the role of drinking goal selection. *J Subst Abuse Treat*. 2006;31:419-24. [PMID: 17084796]
75. Floyd RL, Sobell M, Velasquez MM, Ingersoll K, Nettleman M, Sobell L, et al; Project CHOICES Efficacy Study Group. Preventing alcohol-exposed pregnancies: a randomized controlled trial. *Am J Prev Med*. 2007;32:1-10. [PMID: 17218187]
76. Flynn HA, Marcus SM, Barry KL, Blow FC. Rates and correlates of alcohol use among pregnant women in obstetrics clinics. *Alcohol Clin Exp Res*. 2003;27:81-7. [PMID: 12544010]
77. Grant TM, Ernst CC, Streissguth A, Stark K. Preventing alcohol and drug exposed births in Washington state: intervention findings from three parent-child assistance program sites. *Am J Drug Alcohol Abuse*. 2005;31:471-90. [PMID: 16161730]
78. Meberg A, Halvorsen B, Holter B, Ek IJ, Askeland A, Gaaserud W, et al. Moderate alcohol consumption—need for intervention programs in pregnancy? *Acta Obstet Gynecol Scand*. 1986;65:861-4. [PMID: 3825527]
79. O'Connor MJ, Whaley SE. Brief intervention for alcohol use by pregnant women. *Am J Public Health*. 2007;97:252-8. [PMID: 17194863]
80. Reynolds KD, Coombs DW, Lowe JB, Peterson PL, Gayoso E. Evaluation of a self-help program to reduce alcohol consumption among pregnant women. *Int J Addict*. 1995;30:427-43. [PMID: 7607777]
81. Centers for Disease Control and Prevention (CDC). Motivational intervention to reduce alcohol-exposed pregnancies—Florida, Texas, and Virginia, 1997-2001. *MMWR Morb Mortal Wkly Rep*. 2003;52:441-4. [PMID: 12807086]
82. Halmesmaki E. Alcohol counselling of 85 pregnant problem drinkers: effect on drinking and fetal outcome. *Br J Obstet Gynaecol*. 1988;95:243-7. [PMID: 3370196]
83. Waterson EJ, Murray-Lyon IM. Preventing alcohol related birth damage: a review. *Soc Sci Med*. 1990;30:349-64. [PMID: 2408151]
84. Handmaker NS, Miller WR, Manicke M. Findings of a pilot study of motivational interviewing with pregnant drinkers. *J Stud Alcohol*. 1999;60:285-7. [PMID: 10091968]
85. Saitz R. Alcohol screening and brief intervention in primary care: Absence of evidence for efficacy in people with dependence or very heavy drinking. *Drug Alcohol Rev*. 2010;29:631-40. [PMID: 20973848]
86. Miller WR, Walters ST, Bennett ME. How effective is alcoholism treatment in the United States? *J Stud Alcohol*. 2001;62:211-20. [PMID: 11327187]
87. Babor TF, Steinberg K, Anton R, Del Boca F. Talk is cheap: measuring drinking outcomes in clinical trials. *J Stud Alcohol*. 2000;61:55-63. [PMID: 10627097]
88. Del Boca FK, Noll JA. Truth or consequences: the validity of self-report data in health services research on addictions. *Addiction*. 2000;95 Suppl 3:S347-60. [PMID: 11132362]
89. McCambridge J, Kypri K. Can simply answering research questions change behaviour? Systematic review and meta analyses of brief alcohol intervention trials. *PLoS One*. 2011;6:e23748. [PMID: 21998626]

Current Author Addresses: Dr. Jonas: University of North Carolina at Chapel Hill, Department of Medicine, 5034 Old Clinic Building, CB 7110, Chapel Hill, NC 27599.

Dr. Garbutt: University of North Carolina at Chapel Hill, Department of Psychiatry, CB 7160, Chapel Hill, NC 27599.

Ms. Amick, Ms. Wilkins, and Dr. Harris: Cecil G. Sheps Center for Health Services Research, University of North Carolina at Chapel Hill, 725 Martin Luther King Boulevard, CB 7590, Chapel Hill, NC 27599.

Dr. Brown: 6209 West Sugar Pine Trail, Tucson, AZ 85743.

Dr. Brownley: University of North Carolina at Chapel Hill, Department of Psychiatry, CB 7175, Chapel Hill, NC 27599.

Ms. Council and Ms. Evans: RTI International, 3040 Cornwallis Road, Box 12194, Research Triangle Park, NC 27709.

Dr. Viera: University of North Carolina School of Medicine, Department of Family Medicine, 590 Manning Drive, CB 7595, Chapel Hill, NC 27599.

Mr. Schwartz: University of North Carolina School of Medicine, 1001 Bondurant Hall, CB 9535, Chapel Hill, NC 27599.

Ms. Richmond: 229 Cherrywood Avenue, San Leandro, CA 94577.

Mr. Yeatts: 444 South Blount Street 326, Raleigh, NC 27601.

Ms. Wood: 403 Pritchard Avenue, Chapel Hill, NC 27516.

Author Contributions: Conception and design: D.E. Jonas, J.C. Garbutt, H.R. Amick, K.A. Brownley, C.L. Council, E.M. Richmond, R.P. Harris.

Analysis and interpretation of the data: D.E. Jonas, J.C. Garbutt, H.R. Amick, J.M. Brown, K.A. Brownley, C.L. Council, A.J. Viera, T.M. Wilkins, C.J. Schwartz, E.M. Richmond, J. Yeatts, T.S. Evans, S.D. Wood, R.P. Harris.

Drafting of the article: D.E. Jonas, J.C. Garbutt, H.R. Amick, J.M. Brown, K.A. Brownley, C.L. Council, R.P. Harris.

Critical revision of the article for important intellectual content: D.E. Jonas, J.C. Garbutt, H.R. Amick, C.L. Council, A.J. Viera, R.P. Harris. Final approval of the article: D.E. Jonas, H.R. Amick, J.M. Brown, A.J. Viera, J. Yeatts, R.P. Harris.

Provision of study materials or patients: D.E. Jonas.

Statistical expertise: D.E. Jonas.

Obtaining of funding: D.E. Jonas.


Administrative, technical, or logistic support: H.R. Amick.

Collection and assembly of data: D.E. Jonas, H.R. Amick, K.A. Brownley, C.L. Council, A.J. Viera, C.J. Schwartz, E.M. Richmond, J. Yeatts, T.S. Evans, S.D. Wood.

Appendix Table 1. Key Questions for This Systematic Review

1. What is the direct evidence that screening for alcohol misuse followed by a behavioral counseling intervention, with or without referral, leads to reduced morbidity (e.g., alcohol-related morbidity or alcohol-related accidents and injuries), reduced mortality, or changes in other long-term (≥ 6 mo) outcomes (e.g., health care utilization, sick days, costs, legal issues, or employment stability)?
2. How do specific screening modalities compare with one another for detecting alcohol misuse?
3. What adverse effects are associated with screening for alcohol misuse and screening-related assessment?
- 4a. How do behavioral counseling interventions, with or without referral, compare with usual care for improving intermediate outcomes (e.g., change in mean number of drinks per drinking day or heavy drinking episodes) for people with alcohol misuse as identified by screening?
- 4b. How do specific behavioral counseling approaches, with or without referral, compare with one another for improving intermediate outcomes for people with alcohol misuse as identified by screening?
5. What adverse effects are associated with behavioral counseling interventions, with or without referral, for people with alcohol misuse as identified by screening?
6. How do behavioral counseling interventions, with or without referral, compare with one another and with usual care for reducing morbidity (e.g., alcohol-related morbidity or alcohol-related accidents and injuries), reducing mortality, or changing other long-term (≥ 6 mo) outcomes (e.g., health care utilization, sick days, costs, legal issues, or employment stability) for people with alcohol misuse as identified by screening?
7. To what extent do health care system influences promote or hinder effective screening and interventions for alcohol misuse?

Appendix Figure 1. Analytic framework for screening, behavioral counseling, and referral in primary care to reduce alcohol misuse.




KQ = key question.

Appendix Table 2. Definitions of the Grades of Overall Strength of Evidence*

| Grade | Definition |
|--------------|--|
| High | High confidence that the evidence reflects the true effect. Further research is very unlikely to change our confidence in the estimate of effect. |
| Moderate | Moderate confidence that the evidence reflects the true effect. Further research may change our confidence in the estimate of the effect and may change the estimate. |
| Low | Low confidence that the evidence reflects the true effect. Further research is likely to change our confidence in the estimate of the effect and is likely to change the estimate. |
| Insufficient | Evidence either is unavailable or does not permit estimation of an effect. |

* From reference 28.

Appendix Figure 2. Summary of evidence search and selection.



Appendix Table 3. Characteristics of Included Trials Comparing Behavioral Counseling Interventions With Control Groups

| Study, Year (Reference) | Participants (Alcohol-Dependent), n (%) | Duration, mo | Country or State | Setting | Mean Age, y* | Women, %* | Nonwhite, %* | Baseline Alcohol Consumption, drinks/wk* | Quality |
|---|---|--------------|----------------------------|--|---------------------|-----------|--------------|--|---------|
| Adults | | | | | | | | | |
| Anderson and Scott, 1992 (29) | 154 (NR) | 12 | United Kingdom | 8 PC group practices | 43–45.1 | 0 | NR | 37.9–38.8 | Fair |
| WHO BISG, 1996 (63) | 1559 (0) | 9 | 8, including United States | Outpatient medical settings | 35.9–36.9 | 19.2 | NR | NR | Fair |
| SIP Trial (55–57) | 408 (30.4) | 12 | Germany | 85 general practices | 35.9–36.8 | 31.9 | NR | 21–25.2 | Fair |
| Curry et al, 2003 (31) | 307 (NR) | 12 | Washington | 23 PCPs in an HMO and an urban clinic | 47 | 35 | 20 | 14.2 | Fair |
| Project TREAT (33, 35–37, 39) | 774 (NR)† | 48 | Wisconsin | 17 community PC practices | NR‡ | 38 | 5.6–11.9 | 18.9–19.1 | Good |
| Fleming et al, 2008 (51), and Wilton et al, 2009 (54) | 235 (NR) | 6 | Wisconsin | 34 obstetric practices | Median, 28 | 100 | 18.3 | 8–8.5 | Good |
| Lock et al, 2006 (52) | 127 (0) | 12 | United Kingdom | General practices | 44.1 | 50 | NR | 23–26.48 | Fair |
| ELM Trial (59–61) | 301 (NR) | 12 | Pennsylvania | 12 PC clinics | 45.6 | 30.2 | 23.3 | 15.5–18.6 | Fair |
| Nokoy et al, 2010 (53) | 117 (13.8–15.3)§ | 6 | Thailand | Rural PC units | 37 | 8.5 | 100 (Thai) | 15.15 | Fair |
| Project Health (42–44) | 530 (2) | 48 | Massachusetts | 4 PC sites (93 clinicians) | 43.5–44.2 | 32.1–38.7 | 4.3–6.6 | 16.6–18.9 | Fair |
| Richmond et al, 1995 (45) | 378 (35)¶ | 12 | Australia | 40 PC practices | 37.7 | 43 | NR | 38.5 | Fair |
| Rubio et al, 2010 (46) | 752 (0) | 12 | Spain | 20 PC centers in Madrid | NR; >70% were 31–40 | 34.7 | NR | 26.90–27.42 | Fair |
| Saitz et al, 2003 (47) | 312 (NR) | 6 | Massachusetts | Urban academic PC practice | 42.2–43.7 | 29–43 | 80–82 | Mean, 5.5–5.6 drinks per drinking day | Fair |
| Scott and Anderson, 1991 (49) | 72 (NR) | 12 | United Kingdom | 8 PC group practices | 44.4–47.2 | 100 | NR | 25.8–26.7 | Fair |
| Freeborn et al, 2000 (58), and Senft et al, 1997 (62) | 516 (0) | 24 | Oregon | 3 PC clinics in an HMO | 41.9–43 | 28.1–31.1 | 17.4–18.7 | 16.5 | Fair |
| Wallace et al, 1988 (50) | 909 (NR) | 12 | United Kingdom | 47 group practices | 41.7–44.6 | 29.1–29.8 | NR | 35.1 (women) and 62.2 (men) | Fair |
| Older adults | | | | | | | | | |
| Fleming et al, 1999 (34), and Mundt et al, 2005 (41) | 158 (0) | 24 | Wisconsin | 24 PC practices | NR; >92% were 65–75 | 33.5 | NR | 15.5–16.6 | Fair |
| Lin et al, 2010 (38), and Moore et al, 2011 (40) | 631 (NR) | 12 | California | PC practices (145 PCPs) | 68.4 | 29 | 13 | 15.2 | Fair |
| Young adults/college students | | | | | | | | | |
| Fleming et al, 2010 (32) | 986 (0) | 12 | United States and Canada | 5 college health clinics | 21 | 50.5–51.3 | 8.1–10.5 | 17.3–17.8 | Good |
| Grossberg et al, 2004 (37) | 226 (NR)¶ | 48 | Wisconsin | 17 community PC practices | NR** | 51 | 14 | 16.2–18.3 | Good¶ |
| Kypri et al, 2007 (64), and Kypri et al, 2008 (65) | 576 (NR) | 12 | New Zealand | University primary health care service | 20.1–20.3 | 52 | NR | NR | Good |
| Kypri et al, 2004 (66) | 104 (NR) | 6 | New Zealand | University student health service | 19.9–20.4 | 50 | NR | NR | Fair |
| Schaus et al, 2009 (48) | 363 (0) | 12 | Florida | College student health center | 20.6 | 52 | 22 | 8.4–9.6 | Fair |
| Pregnant women | | | | | | | | | |
| Chang et al, 1999 (30) | 250 (0 current)†† | About 6 | Massachusetts | Obstetric practices | 30.7 | 100 | 22 | Mean, 0.6–0.9 drinks per drinking day†† | Fair |

ELM = Early Lifestyle Modification; NR = not reported/unclear; PC = primary care; PCP = primary care physician; SIP = Screening and Intervention in Primary Care; TREAT = Trial for Early Alcohol Treatment; WHO BISG = World Health Organization Brief Intervention Study Group.

* When data were not reported for the total sample but were presented for each study group, the range of the means for the various study groups are reported.

† According to a medical record audit, 6 participants received formal alcohol treatment during the 1-y follow-up; those 6 may ultimately have received a diagnosis of alcohol dependence.

‡ Group 1, men: age 18–30 y, 20.2%; 31–40 y, 27.2%; 41–50 y, 23.9%; 51–65 y, 28.8%. Group 1, women: age 18–30 y, 43.5%; 31–40 y, 25.9%; 41–50 y, 15.6%; 51–65 y, 15.0%. Group 2, men: age 18–30 y, 26.0%; 31–40 y, 25.1%; 41–50 y, 21.3%; 51–65 y, 27.7%. Group 2, women: age 18–30 y, 35.7%; 31–40 y, 35.7%; 41–50 y, 18.2%; 51–65 y, 10.5%.

§ Based on an Alcohol Use Disorders Identification Test score >25.

¶ Percentage with moderate physical dependence based on the physical dependence score from the Comprehensive Drinker Profile. The study excluded participants with evidence of severe alcohol dependence (physical dependence score >10) or those with severe levels of alcohol-related problems (Michigan Alcohol Screening Test score >20).

¶¶ In this subgroup analysis of TREAT (33), 226 of 774 enrolled participants were young adults (aged 18 to 30 y).

** Age 18–21 y, 21%; 22–25 y, 37%; and 26–30 y, 47%.

†† None of the 250 participants satisfied *Diagnostic and Statistical Manual of Mental Disorders, Third Edition, Revised*, criteria for current alcohol abuse or dependence at enrollment; however, 40% satisfied criteria for lifetime alcohol abuse or dependence (not reported separately) and 3 participants had been treated for an alcohol problem.

‡‡ While pregnant, including abstainers. The reported mean excluding abstainers was 1.5–2.1 drinks per drinking day while pregnant.

Appendix Table 4. Description of Behavioral Counseling Interventions, by Intervention Intensity

| Population and Type of Intervention | Study, Year (Reference) | Intervention | Delivered By | Delivery Method | Contacts, n | Length of Each Contact |
|---|---|--|--|---------------------|---|--|
| Adults | Richmond et al., 1995 (45) | Group 2: Physician advice and a self-help manual (after assessment) | PCP | In person | 1 | 5 min |
| | WHO BISC, 1996 (63) | Group 1: Advice and an illustrated pamphlet | Various clinic staff | In person | 1 | 5 min |
| | Anderson and Scott, 1992 (29) | Brief advice, feedback about own consumption and norms, and a self-help booklet | PCP | In person | 1 | 10 min |
| | Lock et al., 2006 (52) | Brief advice (drink-less protocol) on standard drink units; recommended consumption levels; benefits of cutting down, tips on reducing consumption, advice on goal-setting, action plan, and self-help booklet/diary | Nurse or PA | In person | 1 | 5–10 min |
| | ELIM Trial (59–61) | Brief advice that emphasized feedback from baseline results and implications for drinking, coupled with advice about a goal to reduce or stop alcohol consumption | Research staff | In person | 1 | 10–15 min |
| | Scott and Anderson, 1991 (49) | Brief advice, feedback about own consumption and norms, and a self-help booklet | PCP | In person | 1 | 10 min |
| | Freeborn et al., 2000 (58), and Senft et al., 1997 (62) | 30-s message from PCP and 15-min session with health counselor immediately after PCP visit | PCP and study health counselor | In person | 1 | About 15 min |
| | WHO BISC, 1996 (63) | Group 2: Brief intervention and 30-page, illustrated problem-solving manual | Various clinic staff | In person | 1 | 15 min |
| | Curry et al., 2003 (31) | Brief motivational message from PCP during regularly scheduled visit, self-help manual, written personalized feedback, and ≤3 outreach phone counseling calls from health educator | PCP and research health educator | In person and phone | Up to 4 | 1–5 min PCP; mean for phone calls, 14 min |
| | Project TREAT (33, 35–37, 39) | Two visits with PCP, 1 mo apart, and a follow-up phone call from the clinic nurse 2 wk after each visit; workbook containing feedback about current health behaviors, review of prevalence of problem drinking, list of adverse effects of alcohol, worksheet on drinking cues, drinking agreement or prescription, and drinking diary cards | PCP and nurse | In person and phone | 4 | 15 min for PCP contacts; NR for phone calls |
| Fleming et al., 2008 (51), and Wilton et al., 2009 (54) | Two visits, each with phone follow-up, and a workbook containing scripted messages with feedback about current health behaviors, prevalence of problem drinking, list of adverse effects of alcohol focused on women and pregnancy, worksheet on drinking cues, drinking agreement in the form of a prescription, and drinking diary cards | Nurse (90%) or obstetrician | In person and phone | 4 | 15 min for in-person contacts; NR for phone calls | |
| Noknoy et al., 2010 (53) | Motivational enhancement protocol: Brief counseling sessions that used a patient-centered interviewing style and considered stages of change | Nurse or PA | In person | 3 | 15 min | |
| Project Health (42–44) | Tailored consultation with a clinician plus follow-up visit, counseling (entailed talking about number of drinks/wk, heavy drinking episodes, or both), patient alcohol consumption information and education materials attached to the patient's chart by research assistants at regular office visit, and receipt of a health booklet at enrollment | PCP | In person | 2 | 5–10 min | |
| Rubio et al., 2010 (46) | Brief advice given by using an intervention workbook containing a review of alcohol-related health effects, a pie chart displaying frequency of types of at-risk drinkers, list of methods for cutting down, a treatment contract, and cognitive behavioral exercises; phone reinforcement by a nurse; and a general health booklet | PCP | In person | 2 | 10–15 min | |
| Extended multicontact | Wallace et al., 1988 (50) | Brief advice, an information booklet ("That's the Limit"), sex-based recommendations for limiting drinking, a drinking diary, and follow-up sessions | PCP | In person | 1 to 5* | NRT |
| | Bischof et al., 2008 (55) | Group 1 (full care) comprised immediate computerized postassessment feedback and multiple sessions of counseling by a psychologist | Trained psychologists from research team | Phone | 4 | Scheduled for 30 min each; mean received, 80.3 min |
| | Richmond et al., 1995 (45) | Group 2 (stepped care) comprised immediate computerized postassessment feedback and a maximum of 3 counseling sessions with a psychologist; sessions were discontinued if patients indicated consumption below study criteria and high self-efficacy to maintain desired behavior | Trained psychologists from research team | Phone | Up to 4 | Scheduled for 30–40 min each |
| | | Group 1 (the Alcoholscreen program) comprised 5 short consultations (introduction, patient education, 3 follow-ups) designed to reduce drinking to recommended limits and included a self-help manual, daily alcohol diary, and personalized patient education and counseling | PCP | In person | 5 | 15–20 min (initial); 5–25 min (follow-ups) |

Continued on following page

Appendix Table 4—Continued

| Population and Type of Intervention | Study, Year (Reference) | Intervention | Delivered By | Delivery Method | Contacts, n | Length of Each Contact |
|--------------------------------------|--|---|------------------------------|------------------------------------|------------------|--|
| Not reported or unknown | ELM Trial (55–57) Saltz et al, 2003 (47) | Motivational enhancement: longer, main initial session; 2 shorter booster sessions; use of empathy and other techniques to enhance motivation; focus on feedback of assessment data and setting alcohol-use goals Providing physicians with positive alcohol screening results and specific recommendations for their patients at a visit | Research staff PCP | In person In person | 3 NR/unknown† | 30–45 min (main); 15–20 min (boosters) NR/unknown† |
| Older adults | | | | | | |
| Brief multicontact | Fleming et al, 1999 (34), and Mundt et al, 2005 (41) | General health booklet plus drinking behavior feedback (workbook), review of problem-drinking prevalence, reasons for drinking, adverse effects of alcohol, drinking cues, a “prescribed” drinking agreement, and drinking diary cards | PCP and nurse | In person and phone | 4 | 10–15 min for PCP contacts; NR for phone calls |
| Extended multicontact | Lin et al, 2010 (38), and Moore et al, 2011 (40) | Personalized risk report and diary for tracking alcohol use; PCP gave oral and written advice in prescription style via an alcohol education booklet, followed by additional feedback and counseling with motivational interviewing from a health educator at weeks 2, 4, and 8 | PCP and health educator | In person and phone | 4 | 15–20 min |
| Young adults/college students | | | | | | |
| Brief | Kypri et al, 2007 (64), and 2008 (65) Kypri et al, 2004 (66) | Web-based assessment and personalized feedback on drinking Web-based assessment and personalized feedback on drinking | Self Self | Computer Computer | 1 1 | 10–15 min 10–15 min |
| Brief multicontact | Fleming et al, 2010 (32) | Two visits 1 month apart with PCP and a follow-up phone call or e-mail from the PCP after each visit, feedback about current behaviors, review of prevalence of high-risk drinking among college students, list of alcohol’s adverse consequences relevant to college students, lists of personal likes and dislikes of drinking, worksheets on drinking cues, BAC level calculator, life goals and alcohol effects, prescription agreement, and drinking diary cards | PCP | In person | 4 | 15 min |
| Extended multicontact | Grossberg et al, 2004 (37) Kypri et al, 2007 (64), and 2004 (66) Schaus et al, 2009 (48) | Subgroup analysis of young adults enrolled in Project TREAT (Fleming et al, 1997 [33]) Web-based assessment and personalized feedback on drinking Motivational intervention sessions that combined patient-centered motivational interviewing and cognitive-behavioral skills training plus a booklet on alcohol prevention | PCP and nurse Self PCP | In person Computer In person | 4 3 2 | 15 min 10–15 min 20 min |
| Pregnant women | | | | | | |
| Extended | Chang et al, 1999 (30) | Review of lifestyle changes made since pregnancy, articulation of drinking goals, identification of circumstances in which participant might be tempted to drink, alternatives to drinking in such situations, take-home manual with tailored notes, and U.S. Surgeon General recommendation | PCP and researcher | In person | 1 | 2-h assessment, 45-min intervention |

BAC = blood alcohol content; ELM = Early Lifestyle Modification; NR = not reported; PA = physician assistant; PCP = physician; PCP = primary care physician; TREAT = Trial for Early Alcohol Treatment; WHO BISG = World Health Organization Brief Intervention Study Group.
 * All participants received an invitation to a 1-mo follow-up; other follow-up was offered at 4, 7, and 10 mo at the discretion of the practitioner.
 † Not reported in the article. Physicians were trained to do “up to 15 minutes,” and the author believes sessions were generally 10–15 min (Wallace P. Personal communication).
 ‡ No particular behavioral intervention was required; the intervention was to provide physicians with positive screening results. On the basis of assessment immediately after the visit, some discussion about drinking was reported for 51% (residents) to 74% (faculty) of visits in the intervention group and 70% (residents) and 51% (faculty) in the control group.