

# Counseling to Prevent Skin Cancer

## Recommendations and Rationale

### U.S. Preventive Services Task Force

This statement summarizes the current U.S. Preventive Services Task Force (USPSTF) recommendation on counseling to prevent skin cancer and the supporting scientific evidence, and updates the 1996 recommendation contained in the *Guide to Clinical Preventive Services, Second Edition*.<sup>1</sup> Explanations of the ratings and of the strength of overall evidence are given in Appendix A and Appendix B, respectively. This recommendation and rationale statement, plus complete information on which this statement is based, including evidence tables and references, are available on the USPSTF Web site <http://www.ahrq.gov/clinic/uspstfix.htm><sup>2</sup> and in print by subscription through the AHRQ Clearinghouse (1-800-358-9295 or E-mail at [ahrqpubs@ahrq.gov](mailto:ahrqpubs@ahrq.gov)) and through the National Guideline Clearinghouse™ ([www.guideline.gov](http://www.guideline.gov)).

## Summary of Recommendation

The U.S. Preventive Services Task Force concludes that the evidence is insufficient to recommend for or against routine counseling by primary care clinicians to prevent skin cancer.

### **I recommendation.**

*The USPSTF found insufficient evidence to determine whether clinician counseling is effective in changing patient behaviors to reduce skin cancer risk. Counseling parents may increase the use of sunscreen for children, but there is little evidence to determine the effects of counseling on other preventive behaviors (such as wearing protective clothing, reducing excessive sun exposure, avoiding sun lamps/tanning beds, or practicing skin self-examination) and little evidence on potential harms.*

## Clinical Considerations

- Using sunscreen has been shown to prevent squamous cell skin cancer. The evidence for the effect of sunscreen use in preventing melanoma, however, is mixed. Sunscreens that block both ultraviolet A (UV-A) and ultraviolet B (UV-B) light may be more effective in preventing squamous cell cancer and its precursors than those that block only UV-B light. However, people who use sunscreen alone could increase their risk for melanoma if they increase the time they spend in the sun.
- UV exposure increases the risk for skin cancer among people with all skin types, but especially fair-skinned people. Those who sunburn readily and tan poorly, namely those with red or blond hair and fair skin that freckles or burns easily, are at highest risk for developing skin cancer and would benefit most from sun protection behaviors. The incidence of melanoma among whites is 20 times higher than it is among blacks; the incidence of melanoma among whites is about 4 times higher than it is among Hispanics.
- Observational studies indicate that intermittent or intense sun exposure is a greater risk factor for melanoma than chronic exposure. These studies support the hypothesis that preventing sunburn, especially in childhood, may reduce the lifetime risk for melanoma.
- Other measures for preventing skin cancer include avoiding direct exposure to midday sun (between the hours of 10:00 AM and 4:00 PM) to reduce exposure to ultraviolet (UV) rays and covering skin exposed to the sun (by wearing protective clothing such as broad-brimmed hats, long-sleeved shirts, long pants, and sunglasses).

---

Corresponding Author: Chair, U.S. Preventive Services Task Force, c/o Project Director, USPSTF, Agency for Healthcare Research and Quality, 540 Gaither Road, Rockville, MD 20850; E-mail: [uspstf@ahrq.gov](mailto:uspstf@ahrq.gov).

- The effects of sunlamps and tanning beds on the risk for melanoma are unclear due to limited study design and conflicting results from retrospective studies.
- Only a single case-control study of skin self-examination has reported a lower risk for melanoma among patients who reported ever examining their skin over 5 years. Although results from this study suggest that skin self-examination may be effective in preventing skin cancer, these results are not definitive.

## Scientific Evidence

### Epidemiology and Clinical Consequences

Melanoma is a leading cause of cancer death in the United States. The lifetime risk for dying of melanoma is 0.36% in white men and 0.21% in white women.<sup>3</sup> Between 1973 and 1995, the age-adjusted incidence of melanoma increased more than 100%, from 5.7 per 100,000 people to 13.3 per 100,000 people. The increase in annual incidence rates is likely due to several factors, including increased sun exposure and possibly earlier detection of melanoma. Although primary prevention efforts have focused on young people, the elderly (especially elderly men) bear a disproportionate burden of morbidity and mortality from melanoma and nonmelanoma skin cancer. Men older than age 65 account for 22% of the newly diagnosed cases of malignant melanoma each year and women in the same age group account for 14%. Basal cell and squamous cell carcinomas are more than 10 times as common as melanoma but account for less morbidity and mortality. Squamous cell cancers, however, may account for 20% of all deaths from skin cancer.

### Effectiveness of Available Interventions

Preventive strategies include reducing sun exposure (eg, by wearing protective clothing and using sunscreen regularly), avoiding sunlamps and tanning equipment, and practicing skin self-examination. There is little direct evidence,

however, that any of these interventions reduce skin cancer morbidity or mortality.

### Reducing Sun Exposure

Avoiding direct sunlight by staying indoors or in the shade or by wearing protective clothing is the most effective measure for reducing exposure to ultraviolet light, but there are no randomized trials of sun avoidance to prevent skin cancer. In numerous observational studies, increased sun exposure in childhood and adolescence is associated with increased risk for non-melanoma skin cancer, which usually occurs in sun-exposed areas such as the face.

Recent studies provide a more complex picture of the relationship between sun exposure and melanoma, however. While melanoma incidence is higher in regions near the equator where ultraviolet exposure is most intense, melanoma often occurs in areas of the body not exposed to the sun. In observational studies, intermittent or intense sun exposure was associated with increased risk for melanoma; chronic exposure was associated with lower risk, as was the ability to tan.<sup>4-7</sup>

### Sunlamp and Tanning Bed Avoidance

Six of 19 case-control studies found a positive association between use of sun lamps and melanoma risk, but most did not adjust for recreational sun exposure or for the dosage and timing of sunlamp exposure.<sup>8</sup> Among 9 studies that examined the duration, frequency, or timing of sunlamp or tanning bed exposure, 4 found a positive association, particularly if the dose of exposure was high and if it caused burning.

### Sunscreen Use

Daily sunscreen use on the hands and face reduced the total incidence of squamous cell cancer in a randomized trial of 1,621 residents in Australia (rate ratio [RR], 0.61; 95% confidence interval [CI], 0.46 to 0.81).<sup>9</sup> Sunscreen had no effect on basal cell cancer. Based on this trial, 140 people would need to use sunscreen daily for 4.5 years to prevent 1 case of squamous cell cancer. An earlier randomized trial demonstrated that sunscreen

use reduced solar keratoses, precursors of squamous cell cancers.<sup>10</sup> There are no direct data about the effect of sunscreen on melanoma incidence. An unblinded randomized trial showed children at high risk for skin cancers who used sunscreen developed fewer nevi than those who did not. Several epidemiologic studies have found higher risk for melanoma among users of sunscreens than among non-users.<sup>11–13</sup> A recent meta-analysis of population-based case-control studies found no effect of sunscreen use on risk for melanoma.<sup>14</sup> The conflicting results may reflect the fact that sunscreen use is more common among fair-skinned persons, who are at higher risk for melanoma, than it is among darker-skinned persons; or, this finding may reflect the fact that sunscreen use could be harmful if it encourages longer stays in the sun without protecting completely against cancer-causing radiation.

### Skin Self-Examination

The only evidence for the effectiveness of skin self-examination comes from a single case-control study.<sup>1,15</sup> After adjustment for other risk factors, skin self-examination was associated with lower incidence of melanoma (odds ratio [OR], 0.66; 95% CI, 0.44 to 0.99) and lower mortality from melanoma (OR, 0.37; 95% CI, 0.16 to 0.84), although the definition of “self-examination” was limited. This study did not provide sufficient evidence that skin self-examination would reduce the incidence of melanoma or improve outcomes of melanoma.

### Effectiveness of Counseling

Community and worksite educational interventions have demonstrated effectiveness for increasing the use of skin protection measures, such as wearing hats and long-sleeve shirts and staying in the shade; however, evidence addressing the effectiveness of clinician counseling to prevent skin cancer is extremely limited. Most studies of counseling have examined intermediate outcomes such as knowledge and attitudes rather than changes in behavior. In a recent survey, 60% of pediatricians said that they usually or always counsel patients about skin protection, but advice to use sunscreen is more common than advice about wearing protective clothing or avoiding the midday sun.<sup>16</sup>

Simple reminders and instructional materials for clinicians can overcome some of the barriers to regular counseling. A randomized trial of a community-based intervention involving 10 towns in New Hampshire suggests that office-based counseling by physicians may be an effective component of a multi-modal program to promote skin protection.<sup>17</sup> The proportion of children utilizing some sun protection increased significantly in the intervention towns (from 78% to 87%) compared to a decrease in the control communities (from 85% to 80%). More parents reported receiving some sun protection information from a clinician in the intervention towns. However, most of the change was due to increased sunscreen use rather than to reduced sun exposure.

### Potential Harms of Skin Protection Behaviors

There are limited data regarding potential harms of counseling or of specific skin protection behaviors. A possible result of skin cancer counseling that focuses on the use of sunscreen can lead to a false sense of security, which might lead to more time in the sun. For example, a randomized trial with young adults found that those who used sunscreen with a high sun protection factor (SPF) stayed longer in the sun than those who used sunscreen with a lower SPF.<sup>18</sup> There has been some concern that use of SPF of 15 results in vitamin D deficiency. However, a randomized trial in people over 40 years of age found that sunscreen use over the summer had no effect on 25-hydroxyvitamin D3 levels. Concerns related to sun avoidance include reduced physical activity levels among children and negative effects on mental health. However, no studies have evaluated the effects of sun protection behaviors on these outcomes.

### Recommendations of Others

The American Cancer Society,<sup>19</sup> the American Academy of Dermatology,<sup>20</sup> the American Medical Association,<sup>21</sup> the American Academy of Pediatrics,<sup>22</sup> the American College of Obstetricians and Gynecologists,<sup>23</sup> and a National Institutes of Health consensus panel<sup>24</sup> all recommend patient education concerning sun avoidance and sunscreen use. The

American Academy of Family Physicians recommends sun protection for all with increased sun exposure.<sup>25</sup> The American College of Preventive Medicine (ACPM) concluded that sun-protective measures (eg, clothing, hats, opaque sunscreens) are probably effective in reducing skin cancer but that the evidence does not support discussion of sunscreen and sun protection with every patient. ACPM concluded that evidence is insufficient to advise patients that chemical sunscreens protect against malignant melanoma and that their use may actually lead to increased risk.<sup>26</sup> Recently, the International Agency for Research on Cancer (IARC), part of the World Health Organization, qualified their recommendation for sunscreen use in ways that address the importance of learning more about potential harms of counseling for sunscreen use as follows:

“Sunscreens probably prevent squamous-cell carcinoma of the skin when used mainly during unintentional sun exposure. No conclusion can be drawn about the cancer-preventive activity of topical use of sunscreens against basal-cell carcinoma and cutaneous melanoma. Use of sunscreens can extend the duration of intentional sun exposure, such as sunbathing. Such an extension may increase the risk for cutaneous melanoma.”<sup>27</sup>

The Task Force on Community Preventive Services found insufficient evidence to determine the effectiveness of a range of population-based interventions to reduce unprotected ultraviolet light exposure and recommended additional research on educational and policy approaches, media campaigns, and both health care setting and community-based interventions.<sup>28</sup>

## References

1. U.S. Preventive Services Task Force. Screening for Skin Cancer. *Guide to Clinical Preventive Services*. 2nd ed. Washington, DC: Office of Disease Prevention and Health Promotion; 1996:141–152.
2. Helfand M, Krages K. *Counseling to Prevent Skin Cancer*. Summary of the Evidence for the U.S. Preventive Services Task Force. Rockville, MD: Department of Health and Human Services, Agency for Healthcare Research and Quality. Available at <http://www.ahrq.gov/clinic/3rduspstf/skacoun/skacounsum.htm>.
3. Ries LA, Kosary CL, Hankey BF, Miller BA, Edwards BK, eds. *SEER Cancer Statistics Review, 1973–95*. Bethesda: National Cancer Institute; 1998.
4. Nelemans PJ, Rampen FH, Ruitter DJ, Verbeek AL. An addition to the controversy on sunlight exposure and melanoma risk: a meta-analytical approach. *J Clin Epidemiol*. 1995;48(11):1331–1342.
5. Elwood M, Jopson J. Melanoma and sun exposure: an overview of published studies. *Int J Cancer*. 1997;73(2):198–203.
6. Walter S, King W, Marrett L. Association of cutaneous malignant melanoma with intermittent exposure to ultraviolet radiation: results of a case-control study in Ontario, Canada. *Int J Epidemiol*. 1999;28(3):418–427.
7. Holly EA, Aston DA, Cress RD, Ahn DK, Kristiansen JJ. Cutaneous melanoma in women. I. Exposure to sunlight, ability to tan, and other risk factors related to ultraviolet light. *Am J Epidemiol*. 1995;141(10):923–33.
8. Swerdlow AJ, Weinstock MA. Do tanning lamps cause melanoma? An epidemiologic assessment. *J Am Acad Dermatol*. 1998;38(1):89–98.
9. Green A, Williams G, Neale R, et al. Daily sunscreen application and beta-carotene supplementation in prevention of basal-cell and squamous-cell carcinomas of the skin: a randomised controlled trial. *Lancet*. 1999;354(9180):723–9.
10. Thompson SC, Jolley D, Marks R. Reduction of solar keratoses by regular sunscreen use. *N Engl J Med*. 1993;329(16):1147–51.
11. Autier P, Dore JF, Schifflers E, et al. Melanoma and use of sunscreens: an EORTC case-control study in Germany, Belgium and France. The EORTC Melanoma Cooperative Group. *Int J Cancer*. 1995; 61(6):749–55.
12. Autier P, Dore JF, Cattaruzza MS, et al. Sunscreen use, wearing clothes, and number of nevi in 6- to 7-year-old European children. European Organization for Research and Treatment of Cancer Melanoma Cooperative Group. *J Natl Cancer Inst*. 1998;90(24): 1873–80.

13. Westerdahl J, Olsson H, Masback A, Ingvar C, Jonsson N. Is the use of sunscreens a risk factor for malignant melanoma? [see comments]. *Melanoma Res.* 1995;5(1):59–65.
14. Huncharek M, Kupelnick B. Use of topical sunscreens and the risk of malignant melanoma: a meta-analysis of 9067 patients from 11 case-control studies. *Am J Public Health.* 2002;92(7):1173–1177.
15. Berwick M, Begg CB, Fine JA, Roush GC, Barnhill RL. Screening for cutaneous melanoma by skin self-examination [see comments]. *J Natl Cancer Inst.* 1996;88(1):17–23.
16. Easton A, Price J, Boehm K, Telljohann S. Sun protection counseling by pediatricians. *Arch Pediatr Adolesc Med.* 1997;151(11):1133–1138.
17. Dietrich AJ, Olson AL, Sox CH, Tosteson TD, Grant-Petersson J. Persistent increase in children's sun protection in a randomized controlled community trial. *Prev Med.* 2000;31(5):569–74.
18. Autier P, Dore JF, Negrier S, et al. Sunscreen use and duration of sun exposure: a double-blind, randomized trial. *J Natl Cancer Inst.* 1999;91(15):1304–1309.
19. American Cancer Society (ACS). Cancer Prevention & Early Detection Facts & Figures 2003. Atlanta, Georgia. Available at: [http://www.cancer.org/docroot/STT/content/STT\\_1x\\_Cancer\\_Prevention\\_Early\\_Detection\\_Facts\\_Figures\\_2003.asp](http://www.cancer.org/docroot/STT/content/STT_1x_Cancer_Prevention_Early_Detection_Facts_Figures_2003.asp) OR at <http://www.cancer.org/downloads/STT/CPED2003PWSecured.pdf>. Accessed March 13, 2003.
20. Lim HW, Cooper K. The health impact of solar radiation and prevention strategies: report of the Environmental Council, American Academy of Dermatology. *J Am Acad Dermatol.* 1999;41(1):81–99.
21. Public Education Announcements for Detection of Skin Cancer. American Medical Association House of Delegates Resolution 406 (I-98). Available at <http://a842.g.akamai.net/f/842/3591/3h/www.ama-assn.org/meetings/public/int1998/reports/refd/rtf/ref406.rtf>. Accessed March 12, 2003.
22. Ultraviolet light: a hazard to children (RE9913). American Academy of Pediatrics, Committee on Environmental Health. *Pediatrics.* 1999;104:328–333.
23. American College of Obstetricians and Gynecologists. Primary and preventive care: periodic assessments. ACOG Committee Opinion 246. Washington, DC: ACOG, 2000.
24. Diagnosis and treatment of early melanoma. NIH Consensus Development Conference, January 27–29, 1992. Vol 10, Number 1.
25. “Sun-Safe Guidelines.” American Academy of Family Physicians. Available at <http://www.aafp.org/afp/20000715/375ph.html>. Accessed March 12, 2003.
26. Ferrini RL, Perlman M, Hill L. Skin protection from ultraviolet light exposure: American College of Preventive Medicine Practice Policy Statement. Available at <http://www.acpm.org/skinprot.htm>. Accessed March 12, 2003.
27. International Agency for Research on Cancer (IARC). Handbooks of Cancer Prevention: Sunscreens. 2001;Vol 5. IARC Press: Lyon, France.
28. Centers for Disease Control and Prevention. Preventing Skin Cancer: Findings of the Task Force on Community Preventive Services on reducing exposure to ultraviolet light. *MMWR* 2003; in press.

## Appendix A U.S. Preventive Services Task Force – Recommendations and Ratings

The Task Force grades its recommendations according to one of 5 classifications (A, B, C, D, I) reflecting the strength of evidence and magnitude of net benefit (benefits minus harms):

- A. The USPSTF strongly recommends that clinicians routinely provide [the service] to eligible patients. *The USPSTF found good evidence that [the service] improves important health outcomes and concludes that benefits substantially outweigh harms.*
- B. The USPSTF recommends that clinicians routinely provide [this service] to eligible patients. *The USPSTF found at least fair evidence that [the service] improves important health outcomes and concludes that benefits outweigh harms.*
- C. The USPSTF makes no recommendation for or against routine provision of [the service]. *The USPSTF found at least fair evidence that [the service] can improve health outcomes but concludes that the balance of benefits and harms is too close to justify a general recommendation.*
- D. The USPSTF recommends against routinely providing [the service] to asymptomatic patients. *The USPSTF found at least fair evidence that [the service] is ineffective or that harms outweigh benefits.*
- I. The USPSTF concludes that the evidence is insufficient to recommend for or against routinely providing [the service]. *Evidence that [the service] is effective is lacking, of poor quality, or conflicting and the balance of benefits and harms cannot be determined.*

## Appendix B U.S. Preventive Services Task Force – Strength of Overall Evidence

The USPSTF grades the quality of the overall evidence for a service on a 3-point scale (good, fair, poor):

**Good:** Evidence includes consistent results from well-designed, well-conducted studies in representative populations that directly assess effects on health outcomes.

**Fair:** Evidence is sufficient to determine effects on health outcomes, but the strength of the evidence is limited by the number, quality, or consistency of the individual studies, generalizability to routine practice, or indirect nature of the evidence on health outcomes.

**Poor:** Evidence is insufficient to assess the effects on health outcomes because of limited number or power of studies, important flaws in their design or conduct, gaps in the chain of evidence, or lack of information on important health outcomes.

### Members of the U.S. Preventive Services Task Force

**Alfred O. Berg, MD, MPH, Chair, USPSTF** (Professor and Chair, Department of Family Medicine, University of Washington, Seattle, WA)

**Janet D. Allan, PhD, RN, Vice-chair, USPSTF** (Dean, School of Nursing, University of Maryland Baltimore, Baltimore, MD)

**Paul Frame, MD** (Tri-County Family Medicine, Cohocton, NY, and Clinical Professor of Family Medicine, University of Rochester, Rochester, NY)

**Charles J. Homer, MD, MPH\*** (Executive Director, National Initiative for Children's Healthcare Quality, Boston, MA)

**Mark S. Johnson, MD, MPH** (Chair, Department of Family Medicine, University of Medicine and Dentistry of New Jersey—New Jersey Medical School, Newark, NJ)

**Jonathan D. Klein, MD, MPH** (Associate Professor, Department of Pediatrics, University of Rochester School of Medicine, Rochester, NY)

**Tracy A. Lieu, MD, MPH\*** (Associate Professor, Department of Ambulatory Care and Prevention, Harvard Medical School, Boston, MA)

**Cynthia D. Mulrow, MD, MSc\*** (Clinical Professor and Director, Department of Medicine, University of Texas Health Science Center, San Antonio, TX)

**C. Tracy Orleans, PhD** (Senior Scientist and Senior Program Officer, The Robert Wood Johnson Foundation, Princeton, NJ)

**Jeffrey F. Peipert, MD, MPH\*** (Director of Research, Women and Infants' Hospital, Providence, RI)

**Nola J. Pender, PhD, RN\*** (Professor Emeritus, University of Michigan, Ann Arbor, MI)

**Albert L. Siu, MD, MSPH** (Professor of Medicine, Chief of Division of General Internal Medicine, Mount Sinai School of Medicine, New York, NY)

**Steven M. Teutsch, MD, MPH** (Senior Director, Outcomes Research and Management, Merck & Company, Inc., West Point, PA)

**Carolyn Westhoff, MD, MSc** (Professor of Obstetrics and Gynecology and Professor of Public Health, Columbia University, New York, NY)

**Steven H. Woolf, MD, MPH** (Professor, Department of Family Practice and Department of Preventive and Community Medicine, Fairfax, VA)

\*Member of the USPSTF at the time this recommendation was finalized.